

Infection Control Micro-Learns

User Guide



About the Micro-Learns

The Project Firstline *Infection Control Micro-Learns* are a series of guided infection control discussions that provide brief, on-the-job educational opportunities. Each micro-learn focuses on a single infection control topic and connects infection control concepts to immediate, practical value. Healthcare workers can easily apply the key points to their daily work and perform the recommended actions to keep germs from spreading.

Using the Micro-Learns

The micro-learns can be incorporated into existing opportunities where groups of healthcare workers gather, such as pre-shift “huddles” or team meetings. The sessions should be led or facilitated by an experienced team member with infection control expertise.

Each micro-learn package includes an adaptable discussion guide for the facilitator and one job aid.



Discussion Guide. The discussion guide is not a script. Facilitators are encouraged to adapt the guide for their audience by incorporating relevant and practical questions and ideas. For instance, facilitators can connect the content to the audience’s job duties, facility-specific cases or issues, resources and points of contact, or other information.



Job Aid. The one-page, visual job aid helps to reinforce the key messages of the micro-learn. Facilitators are encouraged to make the job aid available after the micro-learn session, such as in digital or hard copy form.

Notes for Facilitators

- Before presenting a micro-learn, check the policies and protocols at your facility and adapt the content accordingly.
- Build on your knowledge, experience, and awareness to connect the content to local context or relevant recent events so that your audience can apply the concepts confidently.
- The micro-learns reinforce infection control concepts when risks are observed in patients or in the patient environment, not necessarily in visitors or other staff members.
- Remind your audience that if they see a patient in distress—e.g., with shortness of breath, bleeding, or otherwise at risk of immediate harm—they should respond to the emergency according to facility protocols.

